

The drug "OMARIDIN" for preventive purposes

The most expedient use of the drug in cases with:

- rehabilitation after exhausting illnesses (anorexia, cachexia, after operations on the organs of the digestive tract, tuberculosis, other debilitating infectious diseases and oncopathology).

Hepatitis and fibrosis of the liver,

- atherosclerosis

- diseases of the pancreas,

- the need to build muscle tissue and its functions,

- the need for rapid recovery of damaged tissues, wound healing, etc.,

- kidney diseases (instead of unbalanced types of protein that damage the functions of the kidneys: gelatin, bone broths, offal foods);

- family planning (for men, starting 72 days, for women - 48 days prior to conceptual conception);

- Hypospermia, aspermia, and impotence

- frequent colds (in some cases it may be recommended to inhale a product through an inhaler, in small doses under the supervision of a doctor);

- indications for the strengthening of the immune system, as well as for autoimmune diseases,

- the need to regulate the mass and body structure,

- prophylaxis and in a case of worried heart attacks and stroke,

- burn diseases

- prevention of premature aging,

- a syndrome of fatigue and other diseases.

In all these cases, the therapeutic doses of the drug are recommended, at the level of 1,75 - 2,0 g / kg of body weight per day.

The course duration can be recommended from 3 weeks to 6 months.

At the same time, there are no pathogenetic restrictions on the length of use of the product in the diet.